

Impact of social isolation by Coronavirus disease 2019 in food: a narrative review

Impacto do isolamento social pela Coronavirus disease 2019 na alimentação: uma revisão narrativa

Luísa Harumi MATSUO¹  0000-0003-4825-6729
Camila TURECK¹  0000-0002-5672-1400
Luana Pucci de LIMA¹  0000-0002-3914-2554
Patrícia de Fragas HINNIG¹  0000-0002-9348-8513
Erasmus Benício Santos de Moraes TRINDADE¹  0000-0003-1736-4049
Francisco de Assis Guedes de VASCONCELOS¹  0000-0002-6162-8067

ABSTRACT

The objective of this narrative review, with a systematic survey, was to investigate the impact of social isolation on food (eating habits) during the 2019 Coronavirus disease pandemic. A retrieval of studies published from January 1st, 2019 to June 18th, 2020 was carried out in the PubMed, Web of Science and Embase databases. The screening was structured using the terms corresponding to Coronavirus disease 2019, social isolation and food. All steps were carried out independently by three reviewers. Twelve studies were included, totaling 20,789 people assessed, with a predominance of cross-sectional studies of adult population. The studies reviewed had been carried out in Italy (n=4), Spain (n=2), Poland (n=1), China (n=1), India (n=1), United States of America (n=1), South Africa (n=1) and in a continental multicenter (n=1). The analysis of the studies included showed that most people reported that they did not change their usual diet; in addition, an increase in cooking habits, consumption of fruits and vegetables and a decrease in the consumption of alcoholic beverages was observed. A decrease in fish consumption and an increase in comfort foods were also reported. It was found that food habits changed in times of social isolation. We suggest that new

¹ Universidade Federal de Santa Catarina, Centro de Ciências da Saúde, Programa de Pós-Graduação em Nutrição. *Campus* Universitário Reitor João David Ferreira Lima, R. Eng. Agrônomo Andrei Cristian Ferreira, s/n., Trindade, 88040-900, Florianópolis, SC, Brasil. Correspondence to: F.A.G. VASCONCELOS. E-mail: <f.vasconcelos@ufsc.br>.

Support: *Conselho Nacional Brasileiro de Desenvolvimento Científico e Tecnológico* (CNPq, National Council for Scientific and Technological Development) (Process n. 303233/2019-2).

How to cite this article

Matsuo LH, Tureck C, Lima LP, Hinnig PF, Trindade EBSM, Vasconcelos FAG. Impact of social isolation by Coronavirus disease 2019 in food: a narrative review. *Rev Nutr.* 2021;34:e200211. <https://doi.org/10.1590/1678-9865202134e200211>

investigations be carried out in different socioeconomic and demographic frameworks. We propose to continue the study on this theme, with a systematic review with or without meta-analysis.

Keywords: Coronavirus infections. Diet, food and nutrition. Review. Social isolation.

RESUMO

Trata-se de revisão narrativa com busca sistemática, tendo por objetivo investigar o impacto do isolamento social na alimentação (hábitos alimentares) durante a pandemia da Coronavirus Disease 2019. Foi realizado rastreamento de estudos publicados no período de 1º de janeiro de 2019 a 18 de junho de 2020 nas bases PubMed, Web of Science e Embase. A busca foi estruturada utilizando os unitermos correspondentes a Coronavirus Disease 2019, isolamento social e alimentação. Todas as etapas foram executadas de forma independente por três revisores. Foram incluídos 12 estudos, totalizando 20.789 pessoas pesquisadas, com a predominância de estudos de corte transversal na população adulta. Os estudos analisados foram desenvolvidos na Itália (n=4), Espanha (n=2), Polônia (n=1), China (n=1), Índia (n=1), Estados Unidos da América (n=1), África do Sul (n=1) e multicêntrico continental (n=1). A análise dos estudos incluídos evidenciou que a maioria das pessoas relatou não mudar a dieta habitual, observando-se também o aumento do hábito de cozinhar, consumir frutas, legumes e verduras e a diminuição do consumo de bebidas alcoólicas. Constatou-se diminuição do consumo de peixes e aumento das chamadas comfort foods. Verificou-se que a alimentação sofreu alterações em tempos de isolamento social. Sugerimos que novas pesquisas sejam realizadas em diferentes contextos socioeconômicos e demográficos. Propomos a continuidade do estudo desta temática, com realização de revisão sistemática com ou sem metanálise.

Palavras-chave: Infecções por coronavirus. Alimentos, dieta e nutrição. Revisão. Isolamento social.

INTRODUCTION

In March 2020, Coronavirus disease 2019 (Covid-19), an infectious disease caused by the Acute Severe Respiratory Syndrome 2 (SARS-CoV-2) virus, was characterized by the World Health Organization (WHO) as a pandemic [1,2]. According to WHO data, until August 2, 2020, 17,660,523 million cases of the disease had been confirmed worldwide, and in Brazil, these numbers represented on that date 2,662,485 million cases, with 92,475 thousand deaths [3].

Due to the exponential increase of cases and the absence of measures to immunize the population, non-pharmacological interventions, such as Social Isolation (SI), are indicated to mitigate the transmission of the disease virus. The main objective of social isolation is to contain the virus spread among populations, countries and continents and, to postpone the peak of the epidemic curve, thus reducing the demand for health care and the overload of these services [4]. However, the SI can have an impact on the physical and mental health of individuals and promote changes in life habits related to the practice of physical activity and eating [5]. Food choices may be compromised, mainly due to limited access to food shopping places, which can lead to reduced consumption of fresh food, especially fruits, vegetables and fish, to the detriment of ready-to-eat and ultra-processed foods [6]. In addition, emotional factors in response to routine changes and due to the disease itself, can increase the demand for foods rich in sugars or increase the desire for food [7]. Thus, the objective of this narrative review was to investigate the impact of SI on people's eating habits during the Covid-19 pandemic.

METHODS

Narrative review with systematic survey for studies published from January 1, 2019 to June 18, 2020 in the PubMed, Web of Science and Embase databases using uniterms and with the help of Boolean operators, according to Chart 1, without using filters.

Chart 1 – Systematic search strategy used in the narrative review. *Florianópolis* (SC), Brazil, 2020.

PubMed	
Covid-19	
#1	"Coronavirus" [Mesh]
#2	"Coronavirus Infections" [Mesh]
#3	"Pandemics" [Mesh]
#4	"spike protein, SARS-CoV-2" [Supplementary Concept]
#5	"COVID-19" [Supplementary Concept]
#6	"Disease Outbreaks" [Mesh]
#7	"COVID-19" OR "SARS-CoV-2" OR "covid-19 pandemic" OR "2019-nCoV" OR "COVID-19 virus"
#8	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7
Social Isolation	
#9	"Social Isolation" [Mesh]
#10	"Quarantine" [Mesh]
#11	"Social Isolation" OR "Social Distancing" OR Lockdowns OR Lockdown OR Quarantine OR Quarantines OR "Physical Distance" OR "home confinement" OR home OR confinement*
#12	Distance" OR "home confinement" OR home OR confinement*
#13	#9 OR #10 OR #11 OR #12
Food and eating habits	
#15	"Food" [Mesh]
#16	"Diet, Food, and Nutrition" [Mesh]
#17	"Diet" [Mesh]
#18	"Feeding Behavior" [Mesh]
#19	"Diet, Healthy" [Mesh]
#20	"Nutrients" [Mesh]
#21	"Nutritional Status" [Mesh]
#22	"Life Style" [Mesh]
#23	"Quality of Life" [Mesh]
#24	"food habit" OR "food habits" OR "Healthy eating" OR "eating habits" OR nutrition OR nutrient* OR "Food consumption" OR "dietary behavior" OR "Nutrition Status" OR "eating behavior" OR Lifestyle OR Lifestyles OR "Healthy Lifestyles" OR "Healthy Life Style"
#25	#15 OR #16 OR #17 OR #18 OR #20 OR #21 OR #22 OR #23 OR #24
#26	#8 AND #13 AND #25
Embase	
Covid-19	
1	'spike protein sarscov 2'/exp OR 'Coronavirus infection'/de OR 'coronavirus disease 2019'/exp OR 'pandemic'/exp
Social Isolation	
2	'social isolation'/exp OR 'quarantine'/exp OR 'social distancing'/exp OR 'lockdown'/exp OR 'home'/exp
Food and eating habits	
3	'food'/exp/mj OR 'healthy eating'/exp OR 'feeding behavior'/exp OR 'food intake'/exp OR 'nutrition'/exp OR 'nutrient'/exp OR 'diet'/exp/mj OR 'feeding behavior'/exp OR 'nutritional status'/exp OR 'lifestyle'/exp OR 'healthy diet'/exp/mj OR 'healthy lifestyle'/exp/mj OR 'quality of life'/exp
4	1 AND 2 AND 3
Web of Science	
Covid-19	
#1	Coronavirus OR "Coronavirus Infections" OR Pandemic* OR "COVID-19" OR "Disease Outbreaks" OR "SARS-CoV-2" OR "covid-19 pandemic" OR "2019-nCoV" OR "COVID-19 virus"
Social Isolation	
#2	"Social Isolation" OR Quarantine* OR "Social Distancing" OR Lockdown* OR "Physical Distance" OR "home confinement" OR home OR confinement*
Food and eating habits	
#3	Food OR "Diet, Food, and Nutrition" OR Diet OR "Feeding Behavior" OR "Diet, Healthy" OR Nutrients OR "Life Style" OR "Quality of Life" OR "food habit" OR "food habits" OR "Healthy eating" OR "eating habits" OR nutrition OR nutrient* OR "Food consumption" OR "dietary behavior" OR "Nutrition Status" OR "eating behavior" OR Lifestyle* OR "Healthy Lifestyles" OR "Healthy Life Style"
#4	#1 AND #2 AND #3

Note: 2019-nCoV: Novel Coronavirus 2019; Covid-19: Corona Virus Disease 2019; Exp: Exploded; MeSH: Medical Subject Headings; Mj: Major Focus; SARS-Cov-2: Severe Acute Respiratory Syndrome Coronavirus 2.

The studies captured were transferred to the reference manager Mendeley® to remove duplicates and selected in the Rayyan® program in two stages: reading titles and abstracts and reading the full text. In the selection, the eligibility criteria were used: data collection performed during the isolation period by Covid-19 and that evaluated food and/or eating habits. Review and recommendation studies were excluded.

Data extraction was performed in the Microsoft Excel® 2010 program using a standardized and tested form considering the following variables: authors, year and place of study; methodological design; sample characteristics (gender and age); method used for recruiting the sample; methodology used for data collection (such as online platform, telephone interview); instrument to assess diet; data collection period; type of measure adopted by the country to contain Covid-19 (such as SI); and main results related to food. For the collection, analysis and synthesis of the results related to food in the framework of social isolation, the following aspects were considered: quantity consumed, quality (referring to types of food and/or nutrients consumed), number of meals, cooking habits, forms of preparation, forms of acquisition and food availability.

All steps were carried out independently by three of the authors and compared to verify agreement between them. In the case of any divergence, a fourth reviewer was consulted for final decision.

RESULTS

A total of 439 studies were identified. Out of these, 85 studies were excluded because they were duplicates. A total of 354 were reviewed by title and summary and 21 studies were read in full for eligibility. Twelve studies were included in the narrative review. The selection process is shown in Figure 1.

The Table 1 presents the main characteristics of the studies assessed. Seven studies were carried out on the European continent, four in Italy, two in Spain and one in Poland [8-14]. Two studies were carried out on the Asian continent, one in China and one in India [15,16]. One study was carried out in the American continent, in the United States of America and another in the African continent (South Africa) [17,18]. And only one study corresponded to a multinational survey, involving Europe, North Africa, West Asia and the Americas [19].

Cross-sectional studies were predominant (n=11) and there was only one longitudinal survey [10]. All studies were carried out with populations of both genders; the majority included individuals over 18 years of age [12-15,17-19], two assessed specific populations such as athletes and individuals with diabetes and the longitudinal research assessed obese children and adolescents [10,16,18]. The sample sizes ranged from 41 to 7,514, totaling 20,789 people assessed in the 12 studies. Nine studies presented the type of measure established in the country for the containment of Covid-19, all of which described SI as a condition in which leaving home would occur only for essential purposes.

Data collections were carried out using questionnaires administered on online platforms, with a predominance of Google forms®. To collect data on food, two studies used previously validated questionnaires on frequency and eating behavior and Adherence to the Mediterranean Diet [8,12,13]. The others used adapted instruments or not validated.

In the results related to food, the studies reviewed were grouped into three categories: (a) those that evaluated only eating changes during the SI period compared to the previous period [9,10,12,17-19]; (b) those which evaluated food intake only in the SI period [16]; and (c) those which assessed both [8,11,13-15].




Figure 1 – Flowchart for selecting the 12 studies included in the narrative review.

Regarding changes in food eating during the SI shown in Table 2, the following changes were observed in the aspects: amount of food eaten [11,12,14,17,19]; number of meals [8,10,17,19]; cooking habits [8,12,14]; preparation methods [12]; acquisition [8]; food availability [11,12]; general quality of food [9,18]; and eating-related behaviors [17]. With the exception of Gallè *et al.* [9], Nachimuthu *et al.* [16], and Zachary *et al.* [17], all other studies analyzed the change in the consumption of specific drinks and foods, such as sugary drinks, vegetables, fruits, fish and sweets

The proportion of people who reported an increase in the amount of food eaten during SI ranged between 36.3% and 52.9% [11,12,14,17]. One study identified an increase in the mean score of eating out of control (1.80±0.02 to 2.04±0.03) [19].

In the analysis of the number of meals, it is observed that during quarantine, three and four main meals per day (30.3% and 39.3%, respectively) and one and two daily snacks (28.3% and 36,1% respectively) were more prevalent [14]. In addition, 65.0% reported having breakfast daily during the pandemic [14]. In the Ammar *et al.* [19] multicentric study and in the study with obese children and adolescents by Pietrobelli *et al.* [10] there was an increase in the number of meals and snacks throughout the day. Zachary *et al.* [17] observed that in 11.5%, 22.0% and 38.7% of the participants there was an increase in food intake at breakfast, lunch and dinner meals, respectively.

Table 1 – General characteristics of the 12 studies included in the narrative review. *Florianópolis* (SC), Brazil, 2020.

1 of 3

Variables							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Di Renzo <i>et al.</i> [8].	Italy	Cross-sectional	n=3,533 (≥12 years of age)	M: 23.9% F: 76.1%	18-65 years of age: 91.5%	Google Forms	April 5-24, 2020
Food evaluation							
(a) PREDIMED Mediterranean Diet Adherence Screener; (b) Questions about changes in the consumption of specific foods and/or drinks; (c) Questions about the frequency of consumption of specific foods and/or drinks; (d) Questions about meals (number and/or frequency); (e) Questions about food purchase; (f) Perception of diet change.							
Conclusion							
Greater adherence to the MD in the population between 18 and 30 years of age when compared to the younger and older population. Perception of weight gain was observed in 48.6% of the population, while a slight increase in physical activity has been reported in 38.3% of respondents. In addition, 15.0% of respondents turned to farmers or organic purchasing groups for fruits and vegetables, especially in Northern and Central Italy, where BMI values were lower.							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Gallè <i>et al.</i> [9]	Italy	Cross-sectional	n=2,125 (undergraduates)	M: 37.2% F: 62.8%	Middle age: 22.5	Online questionnaire	Last 2 weeks of March 2020
Food evaluation							
(a) Perception of diet change							
Conclusion							
Social isolation did not change the diet and smoking habits, but it did cause a decrease in the level of physical activity. Preventive interventions should also turn restrictive measures into an opportunity to improve lifestyle							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Pietrobelli <i>et al.</i> [10]	Italy	Longitudinal	n=41 (obese children and adolescents)	M: 53.7% F: 46.3%	6-18 years of age	Baseline: Personal interview Lockdown: Telephone interview	Baseline: May 13 to July 30, 2019 Lockdown: March 10 to April 2020
Food evaluation							
(a) Questions about changes in the consumption of specific foods and/or drinks; (b) Questions about meals (number and/or frequency).							
Conclusion							
In children and adolescents, social isolation seems to create an unfavorable environment for the maintenance of healthy lifestyle behaviors. Depending on the duration, the undesirable effects of social isolation can have a lasting impact on the level of adiposity.							
Author	Country	Design	Population	Gender	Age	Data collection method and	Period
Scarmozzino e Visioli [11]	Italy	Cross-sectional	n=1,929	M: 67.0% F: 32.9%	21-65 years of age 84.1%	Google Forms	April 3-15, 2020
Food evaluation							
(a) Questions about frequency of consumption of specific foods and beverages; (b) Questions about changes in the consumption of specific foods and/or drinks.							
Conclusion							
Almost half (49.6%) of the respondents did not change their diet; however, 46.1% of them reported that they were eating more. There was an increase in the consumption of "comfort foods", mainly chocolate, ice cream and desserts (42.5%) and snacks (23.5%). 21.2% of respondents increased their consumption of fresh fruits and vegetables. Purchases of ready meals reduced by almost 50%.							

Table 1 – General characteristics of the 12 studies included in the narrative review. *Florianópolis* (SC), Brazil, 2020.

Variables							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Rodríguez-Pérez <i>et al.</i> [12]	Spain	Cross-sectional	n=7,514 (>18 years of age)	M: 29.3% F: 70.6%	21-65 years of age: 91.3%	Google Forms	March 2020 (during 3 weeks)
Food evaluation							
(a) PREDIMED Mediterranean diet Adherence Screener (validated); (b) Questions about changes in eating and/or cooking habits; (c) Questions about changes in consumption of specific foods and beverages.							
Conclusion							
Improvement in eating behaviors was observed during social isolation, reflected by greater adherence to MedDiet. Health-related food choices included greater intake of fruits, vegetables or legumes, and less consumption of red meat, alcohol, fried foods or sweets compared to usual habits.							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Romeo-Arroyo <i>et al.</i> [13]	Spain	Cross-sectional	n=600 (NI)	F: 50.1%	18-68 years of age	Online survey;	Last week of April
Food evaluation							
(a) Food Groups Consumption Frequency Questionnaire (QFCGA) (validated); (b) Dutch Eating Behavior Questionnaire, Spanish version (validated); (c) Questions about eating and cooking habits; (d) Questions about changes in the consumption of specific foods and/or drinks; (e) List of new habits acquired.							
Conclusion							
Different attitudes were present in Spanish homes, some of them related to low emotional status and possible less healthy eating habits, and others focused on trying to maintain healthier habits.							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Sidor and Rzymiski [14]	Poland	Cross-sectional	n=1,097 (≥18 years of age)	M: 4.9% F: 95.1%	18-25 years of age: 53.6%	Online survey	April 17/ May 1, 2020
Food evaluation							
(a) Questions about changes in eating and/or cooking habits; (b) Questions about the frequency of consumption of specific foods and/or drinks; (c) Questions about meals (number and/or frequency); (d) Level of concern of contracting SARS-CoV-2 during shopping or contact with food.							
Conclusion							
Social isolation can affect eating habits and behaviors (eating more, more snacks, increased alcohol consumption (Poland) and weight change). Overweight and obese individuals were more prone to these changes (frequency of consumption: <vegetables, fruits and legumes and >starchy foods, meat and dairy products).							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Zhao <i>et al.</i> [15]	China	Cross-sectional	n=1,938 (18-80 years of age)	M: 34.3%; F: 65.7%	18-45 years of age: 83.6%	<i>Wenjuanxing</i>	March 2020
Food evaluation							
(a) Family Diet Diversity Index; (b) Questions about food purchase; (c) Questions about consumption of specific foods in the last 24 hours; (d) Questions about specific eating behaviors to deal with Covid-19.							
Conclusion							
Good general food diversity. Less food diversity in areas with a high number of Covid-19 confirmed cases. Eating behaviors to deal with Covid-19 (consumption of vitamin C, probiotics, other dietary supplements, alcohol and vinegar) were associated with greater diversity. The main ways of obtaining food were internal storage and personal purchases.							
Author	Country	Design	Population	Gender	Age	Data collection method	Period
Nachimuthu <i>et al.</i> [16]	India	Cross-sectional	n=100 individuals with diabetes	M: 54 F: 46	≥18-64: 57.0%	Survey Monkey	April 1-15, 2020
Food evaluation							
(a) Question about diet and exercise during lockdown							

Table 1 – General characteristics of the 12 studies included in the narrative review. *Florianópolis (SC), Brazil, 2020.*

3 of 3

Conclusion

Most people investigated reported that they were able to maintain food care as well as being more physically active at home during social isolation.

Author	Country	Design	Population	Gender	Age	Data collection method	Period
Zachary <i>et al.</i> [17]	United States	Cross-sectional	n=173 (>18 years of age)	M: 44.5% F: 55.5%	Average age: 28.1	Survey Monkey	NI

Food evaluation

Weight and Lifestyle Inventory

Conclusion

Approximately 22% of the sample reported weight gain during social isolation. The risk factors were inadequate sleep, snacks after dinner, lack of food restriction, food in response to stress and reduced physical activity.

Author	Country	Design	Population	Gender	Age	Data collection method	Period
Pillay <i>et al.</i> [18]	South Africa	Cross-sectional	n=692 (athletes)	M: 67.0% F: 33.0%	NI	Google Forms	April 28-30, 2020

Food evaluation

(a) Perception of diet change; (b) Questions about specific eating behaviors to deal with Covid-19.

Conclusion

Covid-19 has significant physical and mental effects on athletes, including worsening of nutrition. Medical, nutritional and psychological support is recommended during and after social isolation.

Author	Country	Design	Population	Gender	Age	Data collection method	Period
Ammar <i>et al.</i> [19]	Europe, Africa, Asia and the Americas	Cross-sectional	n=1,047 (≥18 years of age)	M: 46.2% F: 53.8%	18-35 years of age: 55.1%	Google Forms	April 6-11, 2020

Food evaluation

Short Diet Behaviors Questionnaire for Lockdowns

Conclusion

Interim results of the investigation indicate a negative effect of home confinement on physical activity and diet behavior with a significant increase in sitting time and an unhealthy diet, indicative of a more sedentary lifestyle.

Note: BMI: Body Mass Index; Covid-19: Corona Virus Disease 2019; F: Female; M: Male; MD: Mediterranean Diet; MedDiet: Mediterranean Diet; NI: Not Informed; Covid-19: Coronavirus Disease 2019; PREDIMED: PREvención con Dieta MEDiterránea; SARS-Cov-2: Severe Acute Respiratory Syndrome Coronavirus 2.

In two studies, significant increases in cooking habits were observed, reported by 45.7% and 62.3% of those investigated [11,12]. The preparation of homemade recipes was more present in the food routine during isolation, just as there was a decrease in the proportion of participants who reported the consumption of fried foods and a decrease in the proportion of food delivery services use [8]. The most frequently used form of food purchase was supermarket shopping [8,17].

The difficulty in finding food was reported by 27.0% of the sample in the Rodríguez-Pérez *et al.* study [12]. This aspect negatively influenced consumption of fruits and vegetables, as, according to the study by Scarmozzino and Visioli [11], 33.5% of people reported reducing their consumption due to the difficulty in finding them. This aspect has also been reported as a reason for dietary changes [11].

Table 2 – Changes in eating habits due to social isolation observed in the 12 studies included in the narrative review. *Florianópolis (SC), Brazil, 2020.*

Aspects assessed	Variables			Score
	⇌ %	↑ %	↓ %	
Quantity of food intake				
Scarmozzino and Visioli <i>et al.</i> [11]	47.1	52.9	-	-
Rodríguez-Pérez <i>et al.</i> [12]	63.7	36.3	-	-
Sidor and Rzymiski <i>et al.</i> [14]	-	43.5	-	-
Zachary <i>et al.</i> [17]	61.3	38.7	-	-
Ammar <i>et al.</i> [19]	-	-	-	↑
Number of daily meals				
Di Renzo <i>et al.</i> [8]	57.8	23.5	17.5	-
Pietrobelli <i>et al.</i> [10]	-	-	-	↑
Zachary <i>et al.</i> [17]	34.7	65.3	-	-
Ammar <i>et al.</i> [19]	-	-	-	↑
Cooking				
Di Renzo <i>et al.</i> [8]	-	↑	-	-
Rodríguez-Pérez <i>et al.</i> [12]	50.7	45.7	3.6	-
Sidor and Rzymiski <i>et al.</i> [14]	-	62.3	-	-
Preparation (frying)				
Rodríguez-Pérez <i>et al.</i> [12]	73.4	6.3	20.3	-
Acquisition (food delivery)				
Di Renzo <i>et al.</i> [8]	-	-	↓	-
Food availability				
Scarmozzino and Visioli <i>et al.</i> [11]	-	-	33.5	-
Rodríguez-Pérez <i>et al.</i> [12]	-	-	27.0	-
Food quality				
Gallè <i>et al.</i> [9]	41.9	41.0	17.1	-
Pillay <i>et al.</i> [18]	-	53.0	47.0	-

Note: ⇌ Maintenance; ↑ Increase; ↓ Decrease.

In assessing the general quality of the diet, Gallè *et al.* [9] and Pillay *et al.* [18] found improvement in 41.0% and 53.0% of the population, respectively. The quality deterioration was reported by 17.1% and 47.0%, respectively [9,18]. In the study by Nachimuthu *et al.* [16], which included people with diabetes *mellitus*, 80% reported being regularly following a diet and workout routine during SI and 19.0% reported difficulty due to lack of resources [16].

Zachary *et al.* [17] evaluated behaviors related to the stimulus for food. The authors identified that the behaviors reported in the greatest proportion were the desire to eat certain foods (73%) and boredom (73%), followed by stimulation in response to sight and smell (65%), after dinner (65%), eating with friends and family (59%) and in response to stress (52.0%).

The use of food supplements as a behavior related to Covid-19 was mentioned in two studies [15,18]. Pillay *et al.* [18] found that 32.0% of the athletes evaluated reported using supplements during this period to improve immunity. Among these individuals, 36.0% reported the use of multivitamins, 36.0% vitamin C, 15.0% protein, 4.0% zinc, 23.0% reported other supplements and 5% provided no information [18]. In the study by Zhao *et al.* [15], 37.7% of the interviewees reported making intentional use of dietary supplements, Chinese herbs or specific foods due to Covid-19, including vitamin C (18.2%), probiotics (11.7%), other food supplements (8.0%) and Chinese herbs (9.6%). A few studies have also assessed the change in intake of specific food and beverage groups [8,10-15,18,19], presented in Table 3.

Table 3 – Changes in eating habits due to social isolation observed in the 12 studies included in the narrative review. *Florianópolis* (SC), Brazil, 2020.

Aspects assessed	Variables		
	↔ %	↑ %	↓ %
Fruits and vegetables			
Di Renzo <i>et al.</i> [8]	-	37.4	35.8
Scarmozzino and Visioli <i>et al.</i> [11]	69.2	21.2	8.7
Romeo-Arroyo <i>et al.</i> [13]	-	>30.0	-
Fruits and vegetables (canned)			
Scarmozzino and Visioli <i>et al.</i> [11]	-	0.9	-
Fish¹			
Di Renzo <i>et al.</i> [8]	-	-	-
Scarmozzino and Visioli <i>et al.</i> [11]	64.8	8.5	13.7
Romeo-Arroyo <i>et al.</i> [13]	-	-	>33.0
Meat			
Pietrobelli <i>et al.</i> [10]	-	-	↑
Scarmozzino and Visioli <i>et al.</i> [11]	75.1	4.0	14.9 ²
Romeo-Arroyo <i>et al.</i> [13]	-	>20.0	-
Processed Meat			
Scarmozzino and Visioli <i>et al.</i> [11]	-	6.0	-
Milk/yogurt			
Scarmozzino and Visioli <i>et al.</i> [11]	76.3	14.3	9.4
Cheese			
Scarmozzino and Visioli <i>et al.</i> [11]	79.1	13.3	7.6
Pasta			
Romeo-Arroyo <i>et al.</i> [13]	-	~40.0	-
Olive oil			
Scarmozzino and Visioli <i>et al.</i> [11]	93.2	5.0	1.8
Snacks			
Scarmozzino and Visioli <i>et al.</i> [11]	57.6	23.5	18.9
Rodríguez-Pérez <i>et al.</i> [12]	46.7	37.6	15.7
Fast-food			
Rodríguez-Pérez <i>et al.</i> [12]	60.0	5.1	34.9
Ready meals			
Scarmozzino and Visioli <i>et al.</i> [11]	47.4	3.1	49.5
Junk food			
Di Renzo <i>et al.</i> [8]	-	25.6	29.8
Ammar <i>et al.</i> [19]	-	↑	-
Potato chips			
Pietrobelli <i>et al.</i> [10]	-	↑	-
Crackers			
Scarmozzino and Visioli <i>et al.</i> [11]	66.1	19.1	14.8
Sweets			
Di Renzo <i>et al.</i> [8]	-	-	↓
Scarmozzino and Visioli <i>et al.</i> [11]	44.0	42.5	13.5
Romeo-Arroyo <i>et al.</i> [13]	-	>50.0	-
Sugary Drinks			
Pietrobelli <i>et al.</i> [10]	-	↑	-
Scarmozzino and Visioli <i>et al.</i> [11]	72.7	5.0	22.3
Coffee/tea/infusions			
Di Renzo <i>et al.</i> [8]	-	↑	-
Scarmozzino and Visioli <i>et al.</i> [11]	58.7	29.8	11.5
Alcohol			
Di Renzo <i>et al.</i> [8]	-	-	↓
Scarmozzino and Visioli <i>et al.</i> [11]	53.1	10.1	36.8
Rodríguez-Pérez <i>et al.</i> [12]	32.3	10.4	57.3
Romeo-Arroyo <i>et al.</i> [13]	-	>20.0	-
Sidor and Rzymyski <i>et al.</i> [14]	77.0	14.6	-
Zhao <i>et al.</i> [15]	-	10.6	-
Pillay <i>et al.</i> [18]	-	10.0	-
Ammar <i>et al.</i> [19]	-	-	↓

Note: ↔ Maintenance; ↑ Increase; ↓ Decrease; ¹Fresh and canned fish; ²Red and processed meats.

The consumption of alcoholic beverages was assessed in eight studies [8,11-15,18,19]. Rodríguez-Pérez *et al.* [12] found a predominance in the decrease and maintenance of alcohol consumption, reported by 57.3% and 32.3% of the sample, respectively. Ammar *et al.* [19] also observed a reduction in the mean alcohol binge score (1.15 ± 0.01 to 1.08 ± 0.01) [19]. On the other hand, in the studies by Scarmozzino and Visioli [11] and Sidor and Rzymiski [14], there was a higher proportion of reports indicating maintenance of the usual consumption of alcoholic beverages during the isolation period, representing 53.1% and 77.0% of the samples evaluated, respectively.

The study by Scarmozzino and Visioli [11] found that the majority of the population maintained the usual consumption of sugary drinks (juices, soft drinks, tonic, etc.) (72.7%), with 79.8% of people consuming less than once a week [11]. In contrast, in Pietrobelli *et al.* [10], the longitudinal analysis with children and adolescents identified an increase in the average daily frequency of consumption during the SI (0.90 ± 1.16) when compared to the previous period (0.40 ± 0.90) [10]. Pillay *et al.* [18] mentioned that among the athletes who reported worsening of the diet during the SI (53% of the total sample), 30.0% reported to be due to the consumption of carbonated drinks. The increase in the consumption of hot drinks (coffee, tea and infusions) was reported by more than 20% of the individuals studied by Di Renzo *et al.* [8] and was also mentioned by 29.8% of the population in the study by Scarmozzino and Visioli [11].

Maintaining the consumption of pasta, rice, bread and cereals was mentioned by more than 50% of the individuals in the studies by Romeo-Arroyo *et al.* [13]. In contrast, Di Renzo *et al.* [8] observed that 20% of the study population increased cereal consumption.

Most individuals maintained their consumption of legumes according to the study by Romeo-Arroyo *et al.* [13], but there was an increase in legumes consumption by more than 10% of the study individuals, according to Di Renzo *et al.* [8]. The prevalence of daily consumption of legumes was 70.1% according to Zhao *et al.* [15]. Although it was found that the majority of individuals maintained their consumption of fruits and/or vegetables during SI, it is highlighted that 21.2% and more than 30% of the population reported a consumption increase of these foods [11,13].

Romeo-Arroyo, *et al.* [13] highlighted that the majority of individuals maintained the consumption of meat and eggs. However, Pietrobelli *et al.* [10] observed a significant increase ($p<0.05$) in the average consumption of red meat in children and adolescents as compared to the previous period (1.80 ± 1.53 servings/day) and during the SI (3.46 ± 2.45 servings/day) [10]. Still, among those who mentioned worsening of the diet in the study by Pillay *et al.* [18], 15% of the athletes reported that their response was on account of the excess consumption of red meat.

Di Renzo *et al.* [8] highlighted an increase in the consumption of white meat (not including fish) in more than 10% of people. A decrease in fish consumption was found in several studies: Di Renzo *et al.* [8] (reduction in more than 20% of individuals), Romeo-Arroyo *et al.* [13] (reduction in more than 33.0% of the population) and Scarmozzino and Visioli [11]. In the latter, the authors found that 64.8% of individuals consumed fresh and canned fish as usual, 8.5% increased and 13.7% reduced consumption. In addition, 13.0% of people were consuming less fresh fish and more canned fish [11].

The change in the consumption of milk and dairy products and cheese was evaluated only by Scarmozzino and Visioli [11]. The population studied by them reported usual consumption for these food groups, representing 76.3% for reports of consumption of milk and dairy products and 19.1% for cheese consumption. [11].

Snacks and fast foods showed consumption similar to the period before social isolation, representing 46.7% and 57,6% for snacks, and 60% for fast-foods [11,12]. Additionally, Scarmozzino and Visioli [11]

evaluated the change in the consumption of ready meals and found a higher proportion of reports of maintenance or decrease in consumption, with values of 47.4% and 49.5%, respectively.

In the longitudinal study by Pietrobelli *et al.* [10], children and adolescents reported a significant increase in the consumption of daily servings of potato chips with an average of 0.07 ± 0.24 to 0.61 ± 0.83 . In the study by Ammar *et al.* [19] the average score of frequency of junk food consumption increased significantly during the SI, from 2.18 ± 0.02 to 2.27 ± 0.03 . However, in the study by Di Renzo *et al.* [8] the proportion of increase and decrease reports on the consumption of these foods was similar in the evaluated population, representing 25.6% and 29.8%, respectively.

Scarmozzino and Visioli [11] observed in the studied population a greater predominance of usual consumption of cookies (66.1%). Usual consumption of sweets was reported in 44.0% and the authors observed an increase in 42.5% of the population for this food [11]. Romeo-Arroyo *et al.* [13] found similar values regarding the increase in the consumption of sweets in more than 50% of the individuals assessed.

DISCUSSION

This review included studies that investigated changes in consumption and/or eating habits in times of SI due to Covid-19. We found that in general people maintained the usual diet, although some studies have also found favorable and/or unfavorable changes in the diet.

As favorable changes in eating habits, we can highlight the increase in the habit of cooking and the decrease in the preparation of fried foods. Changes in these aspects may be related to the greater time availability for preparing meals. In a survey of university students ($n=350$), 51% reported that time is an important factor to have a healthy diet, and having time to cook was referred to as being very (47%) and extremely (29%) important for this practice [20].

The increase in the consumption of fruits and vegetables, as well as the greater adherence to the Mediterranean diet, can be acknowledged as a positive change. This observation is in line with the publication of the WHO guidelines on food and nutrition for the isolation period. In this guide, WHO refers to fruits and vegetables as the best food to be consumed in the isolation period [21]. In addition, fruits and vegetables are sources of essential micronutrients for the immune system, mainly vitamins A, C, D, E, B2, B6 and B12, folic acid, iron, selenium and zinc [22]. Another nutrient that favors the immune function and has an anti-inflammatory characteristic is omega-3 fatty acid, found in fish [23]. However, in our study, we identified a decrease in the consumption of this food, which may suggest a difficulty in the ability to prepare this type of food or in the reduction of the purchase at specific places where fish is sold [24].

Among the diet changes considered negative for health, we can mention increase in the consumption of foods with high energy density, high concentration of sugar and fat (sweets, sugary drinks, snacks and sauces), also called comfort foods. The increase in consumption of these foods may be due to the increase in the levels of anxiety and stress and is related with negative health outcomes [25,26].

The results observed in only one study with obese children and adolescents indicate unfavorable changes in the diet, such as an increase in the intake of sugary drinks and potato chips. The habitual consumption of these foods is associated with an increased risk of overweight and obesity [27]. It is noteworthy that the behaviors and habits acquired in this phase of life tend to be perpetuated in adult life, an aspect of great concern in a scenario of continuity of measures of social isolation for an indefinite period [28]. Thus, there is a need for further studies in this population to target interventions during this period.

Although the aim of this review was not the search for articles that exclusively evaluated alcohol consumption during the pandemic, the studies included indicated greater proportions of reduction and maintenance of alcohol consumption. However, it is known that alcohol consumption during the pandemic can represent an important public health problem [29]. During social isolation related to Covid-19, individuals may experience greater anxiety and depressive symptoms and evidence suggests that psychological suffering related to Covid-19 is associated with increased alcohol consumption [30,31]. A study in Poland found that 17.4% of the sample reported a decrease in the consumption of alcoholic beverages due to the pandemic, 13.8% an increased consumption and about 41.1% did not report changes [32].

The decrease in alcohol consumption was observed concomitantly with the increase in the use of coffee/tea/infusions. One hypothesis is that staying at home without contact with other people has influenced the pattern of drink consumed, since alcoholic beverages are related to the socialization process, as a set of strategies that facilitate the coexistence between people, especially young people [33].

We observed that the use of dietary supplements was a strategy used during SI. Some authors and even social media professionals have recommended the use of supplements to enhance immunity [34,35]. Easy access to information may have favored the adoption of this behavior; however supplementation without the guidance of a trained professional is not advisable.

There was an increase in the number of meals among children and adolescents [10]. A study, carried out not in connection with the pandemic, with a population in the same age group, found that a greater number of meals at home may be associated with better quality of the diet and lower intake of saturated fat [36]. This behavior was also observed in studies with a predominantly adult sample [8,17,19]. However, in the SI situation, it is not possible to infer that the increase in the number of meals can be beneficial for the diet, since it can be associated with the consumption of foods considered comfort food [25].

Finally, we highlight that according to the studies included in this review, it is not possible to infer whether food changes differ between countries, since the studies present different recruitment methodologies and data collection on food consumption, as well as the variability of the groups and types of foods assessed. Still, the method of data collection (online questionnaire), necessary due to social isolation, can be an important factor of sample selection bias, including populations with higher education and income, as well as younger populations, depending on the availability and familiarity with digital technologies. Most of the studies found so far have occurred in countries with a high Human Development Index (HDI) (n=8), with limited results in developing countries and no studies in low HDI countries. [37]. We emphasize that Low and Middle-Income (LMIC) countries can suffer an even greater (or different) impact in the isolation process.

The potentialities of this study include: the current and relevant subject when covering this pandemic period and the methodological rigor in the stages of search, selection and data extraction, according to the recommendations of Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA). As limitations, there is the time limit considering the continuity of the pandemic.

We suggest that new studies be carried out in different countries, age groups and economic levels. The studies should also focus on the impact of diet changes in times of isolation on future outcomes, with follow-up studies. We emphasize the importance of developing and validating self-applicable online tools that can reach different population groups, thus providing valid data for assessing food consumption in similar periods in the future. Finally, we propose to continue the study of this topic, with a systematic review with or without meta-analysis.

CONCLUSION

Social isolation can impact changes in the eating habits of different countries' populations. There was an increase in the report of the cooking habits and consumption of fruits and vegetables. On the other hand, a decrease of fish consumption and an increase in comfort foods was reported. The decrease in the consumption of alcoholic beverages should be viewed with caution, since the present review was not limited to articles that exclusively evaluated their consumption. It is suggested that further studies be carried out in different socioeconomic and demographic contexts.

ACKNOWLEDGMENTS

We are grateful to the *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* (CAPES, Coordination of Superior Level Stass Improvement) for their support.

CONTRIBUTORS

LH MATSUO, C TURECK, and LP LIMA contributed to the conception and design of the study, search, collection, analysis and interpretation of data, writing of the manuscript and final version approval. PF HINNIG, EBSM TRINDADE, and FAG VASCONCELOS contributed to the conception and design of the study, analysis and interpretation of data, review and approval of the final version of the article.

REFERENCES

1. World Health Organization. WHO announces COVID-19 outbreak a pandemic 2020. Geneva: Organization; 2020 [cited 2020 Aug 2]. Available from: <https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who-announces-covid-19-outbreak-a-pandemic>
2. Guo Y-R, Cao Q-D, Hong Z-S, Tan Y-Y, Chen S-D, Jin H-J, *et al.* The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak: an update on the status. *Mil Med Res.* 2020;7:11. <https://doi.org/10.1186/s40779-020-00240-0>
3. World Health Organization. Coronavirus Disease (COVID-19) Dashboard 2020. Geneva: Organization; 2020 [cited 2020 Aug 2]. Available from: <https://covid19.who.int/>
4. Ministério da Saúde (Brasil). Medidas não farmacológicas 2020. Brasília: Ministério; 2020 [citado 2 Ago 2020]. Disponível em: <https://coronavirus.saude.gov.br/medidas-nao-farmacologicas>
5. Garcia LP, Duarte E. Intervenções não farmacológicas para o enfrentamento à epidemia da COVID-19 no Brasil. *Epidemiol Serv Saúde.* 2020;29:e2020222. <https://doi.org/10.5123/S1679-49742020000200009>
6. Kanter R, Boza S. Strengthening local food systems in times of concomitant global crises: reflections from Chile. *Am J Public Health.* 2020;110:971-3. <https://doi.org/10.2105/AJPH.2020.305711>
7. Muscogiuri G, Barrea L, Savastano S, Colao A. Nutritional recommendations for CoVID-19 quarantine. *Eur J Clin Nutr.* 2020;74:850-1. <https://doi.org/10.1038/s41430-020-0635-2>
8. Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attinà A, Cinelli G, *et al.* Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med.* 2020;18:229. <https://doi.org/10.1186/s12967-020-02399-5>
9. Gallè F, Sabella EA, Da Molin G, De Giglio O, Caggiano G, Di Onofrio V, *et al.* Understanding knowledge and behaviors related to CoViD-19 epidemic in Italian undergraduate students: the EPICO study. *Int J Environ Res Public Health.* 2020;17(10)3481. <https://doi.org/10.3390/ijerph17103481>
10. Pietrobelli A, Pecoraro L, Ferruzzi A, Heo M, Faith M, Zoller T, *et al.* Effects of covid-19 lockdown on lifestyle behaviors in children with obesity living in Verona, Italy: a longitudinal study. *Obesity.* 2020;28:1382-5. <https://doi.org/10.1002/oby.22861>

11. Scarmozzino F, Visioli F. Covid-19 and the subsequent lockdown modified dietary habits of almost half the population in an Italian sample. *Foods (Basel, Switzerland)*. 2020;9(5):675. <https://doi.org/10.3390/foods9050675>
12. Rodríguez-Pérez C, Molina-Montes E, Verardo V, Artacho R, García-Villanova B, Guerra-Hernández EJ, *et al*. Changes in dietary behaviours during the COVID-19 outbreak confinement in the Spanish COVIDiet study. *Nutrients*. 2020;12(6):1730. <https://doi.org/10.3390/nu12061730>
13. Romeo-Arroyo E, Mora M, Vázquez-Araújo L. Consumer behavior in confinement times: food choice and cooking attitudes in Spain. *Int J Gastron Food Sci*. 2020;21:100226. <https://doi.org/10.1016/j.ijgfs.2020.100226>
14. Sidor A, Rzymiski P. Dietary choices and habits during COVID-19 lockdown: experience from Poland. *Nutrients*. 2020;12(6):1657. <https://doi.org/10.3390/nu12061657>
15. Zhao A, Li Z, Ke Y, Huo S, Ma Y, Zhang Y, *et al*. Dietary diversity among Chinese residents during the COVID-19 outbreak and its associated factors. *Nutrients*. 2020;12(6):1699. <https://doi.org/10.3390/nu12061699>
16. Nachimuthu S, Vijayalakshmi R, Sudha M, Viswanathan V. Coping with diabetes during the COVID-19 lockdown in India: results of an online pilot survey. *Diabetes Metab Syndr Clin Res Rev*. 2020;14:579-82. <https://doi.org/10.1016/j.dsx.2020.04.053>
17. Zachary Z, Brianna F, Brianna L, Garrett P, Jade W, Alyssa D, *et al*. Self-quarantine and weight gain related risk factors during the COVID-19 pandemic. *Obes Res Clin Pract*. 2020;14(3):210-6. <https://doi.org/10.1016/j.orcp.2020.05.004>
18. Pillay L, Janse van Rensburg DCC, Jansen van Rensburg A, Ramagole DA, Holtzhausen L, Dijkstra HP, *et al*. Nowhere to hide: the significant impact of coronavirus disease 2019 (COVID-19) measures on elite and semi-elite South African athletes. *J Sci Med Sport*. 2020;23:670-9. <https://doi.org/10.1016/j.jsams.2020.05.016>
19. Ammar A, Brach M, Trabelsi K, Chtourou H, Boukhris O, Masmoudi L, *et al*. Effects of COVID-19 home confinement on eating behaviour and physical activity: results of the ECLB-COVID19 international online survey. *Nutrients*. 2020;12:1-13. <https://doi.org/10.3390/nu12061583>
20. Maia RP, Recine E. Valores e práticas sobre alimentação de estudantes da universidade de Brasília. *Demetra Aliment Nutr Saúde*. 2015;10:3-25. <https://doi.org/10.12957/demetra.2014.14215>
21. World Health Organization. Food and nutrition tips during self-quarantine 2020. Geneva: Organization; 2020 [cited 2020 July 30]. Available from: <https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/technical-guidance/food-and-nutrition-tips-during-self-quarantine>
22. Maggini S, Pierre A, Calder PC. Immune function and micronutrient requirements change over the life course. *Nutrients*. 2018;10(10):1531. <https://doi.org/10.3390/nu10101531>
23. Saini RK, Keum Y-S. Omega-3 and omega-6 polyunsaturated fatty acids: dietary sources, metabolism, and significance: a review. *Life Sci*. 2018;203:255-67. <https://doi.org/10.1016/j.lfs.2018.04.049>
24. Lopes IG, Oliveira RG, Ramos FM. Perfil do consumo de peixes pela população brasileira. *Biota Amaz*. 2016;6:62-5. <https://doi.org/10.18561/2179-5746/biotaamazonia.v6n2p62-65>
25. Leng G, Adan RAH, Belot M, Brunstrom JM, Graaf K, Dickson SL, *et al*. The determinants of food choice. *Proc Nutr Soc*. 2017;76:316-27. <https://doi.org/10.1017/S002966511600286X>
26. Payab M, Kelishadi R, Qorbani M, Motlagh ME, Ranjbar SH, Ardalan G, *et al*. Association of junk food consumption with high blood pressure and obesity in Iranian children and adolescents: the CASPIAN-IV study. *J Pediatr (Rio J)*. 2015;91:196-205. <https://doi.org/10.1016/j.jped.2014.07.006>
27. Costa CS, Del-Ponte B, Assunção MCF, Santos IS. Consumption of ultra-processed foods and body fat during childhood and adolescence: a systematic review. *Public Health Nutr*. 2018;21:148-59. <https://doi.org/10.1017/S1368980017001331>
28. Llewellyn A, Simmonds M, Owen CG, Woolacott N. Childhood obesity as a predictor of morbidity in adulthood: a systematic review and meta-analysis. *Obes Rev*. 2016;17:56-67. <https://doi.org/10.1111/obr.12316>
29. Clay JM, Parker MO. Alcohol use and misuse during the COVID-19 pandemic: a potential public health crisis? *Lancet Public Health*. 2020;5(5):e259. [https://doi.org/10.1016/S2468-2667\(20\)30088-8](https://doi.org/10.1016/S2468-2667(20)30088-8)
30. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, *et al*. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395:912-20. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
31. Rodriguez LM, Litt DM, Stewart SH. Drinking to cope with the pandemic: the unique associations of COVID-19-related perceived threat and psychological distress to drinking behaviors in American men and women. *Addict Behav*. 2020;110:106532. <https://doi.org/10.1016/j.addbeh.2020.106532>

32. Chodkiewicz J, Talarowska M, Miniszewska J, Nawrocka N, Bilinski P. Alcohol consumption reported during the COVID-19 pandemic: the initial stage. *Int J Environ Res Public Health*. 2020;17:4677. <https://doi.org/10.3390/ijerph17134677>
33. Rosa LFA, Nascimento ARA. Representações sociais de bebida alcoólica para homens universitários. *Arq Bras Psicol*. 2015;67:3-19.
34. Richards G, Mer M, Schleicher G, Stacey S. COVID-19 and the rationale for pharmacotherapy: a South African perspective. *Wits J Clin Med*. 2020;2:11-8. <https://doi.org/10.18772/26180197.2020.v2nSla2>
35. Jukic I, Calleja-Gonzalez J, Cos F, Cuzzolin F, Olmo J, Terrados N, *et al.* Strategies and solutions for team sports athletes in isolation due to COVID-19. *Sports*. 2020;8(4):56. <https://doi.org/10.3390/sports8040056>
36. Prado BG, Hinnig PF, Tanaka LF, Latorre MRDO. Qualidade da dieta de escolares de 7 a 10 anos do município de São Paulo: associação com o número e os locais de refeições. *Rev Nutr*. 2015;28(6):607-18. <https://doi.org/10.1590/1415-52732015000600004>
37. United Nations Development Programme. Human development report 2019. New York; 2019 [cited 2020 July 30]:352. Available from: <http://hdr.undp.org/sites/default/files/hdr2019.pdf>

Receivid: August 10, 2020
Final version: October 10, 2020
Approved: November 27, 2020